

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/00	A2	(11) International Publication Number: WO 00/59489 (43) International Publication Date: 12 October 2000 (12.10.00)
(21) International Application Number: PCT/US00/08707 (22) International Filing Date: 31 March 2000 (31.03.00) (30) Priority Data: 60/127,939 6 April 1999 (06.04.99) US (71) Applicant: SEPRACOR INC. [US/US]; 111 Locke Drive, Marlborough, MA 01752 (US). (72) Inventors: BARBERICH, Timothy, J.; 40 Elm Street, Concord, MA 01742 (US). RUBIN, Paul, D.; 37 Greystone Lane, Sudbury, MA 01776 (US). YELLE, William, E.; 20 Ernie's Drive, Littleton, MA 01460 (US). (74) Agents: INSOGNA, Anthony, M. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHODS AND COMPOSITIONS FOR THE TREATMENT OF NEUROLEPTIC AND RELATED DISORDERS USING ZIPRASIDONE METABOLITES (57) Abstract The invention relates to novel methods using, and pharmaceutical compositions comprising, ziprasidone metabolites. The methods and compositions of the invention are suitable for the treatment of neuroleptic and related disorders. The invention further encompasses methods of preparing ziprasidone sulfoxide and ziprasidone sulfone.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


**METHODS AND COMPOSITIONS FOR
THE TREATMENT OF NEUROLEPTIC AND
RELATED DISORDERS USING ZIPRASIDONE METABOLITES**

1. FIELD OF INVENTION

5 The invention relates to methods of using, and compositions comprising, ziprasidone metabolites.

2. BACKGROUND OF THE INVENTION

10 Ziprasidone, chemically named (5-[2-{4-(1,2-benzisothiazol-3-yl)piperazin-1-yl}ethyl]-6-chlorooxindole)hydrochloride hydrate, is a substituted benzisothiazolylpiperazine. The free base of ziprasidone has the following structure:




20 Ziprasidone and some of its uses are described by U.S. Patent Nos. 4,831,031 and 5,312,925.


 Like clozapine and risperidone, ziprasidone is a highly potent and selective 5-HT₂ receptor and dopamine D₂ receptor antagonist. Seeger, T.F. *et al.*, J. Pharmacol. Exp. Ther., 275(1):101-113 (1995). Ziprasidone is characterized as an antipsychotic, but may also have anxiolytic and antidepressant effects due to its ability to inhibit serotonin and noradrenaline reuptake. Davis, R. and Markham, A., CNS Drugs, 8(2):154-159 (1997). The therapeutic potential of ziprasidone may also be enhanced by its high affinity for the 5-HT_{1A}, 5-HT_{1D}, 5-HT_{2C} receptor subtypes. Seeger, T.F. *et al.*, J. Pharmacol. Exp. Ther., 275(1):101-113 (1995).

 The metabolism of ziprasidone is complex. When administered orally to healthy humans, the drug is extensively metabolized by at least four major pathways: 1) N-dealkylation of the ethyl side chain attached to the piperazinyl nitrogen; 2) oxidation at sulfur resulting in the formation of sulfoxide or sulfone; 3) reductive cleavage of the benzisothiazole moiety; and 4) hydration of the C=N bond and subsequent sulfur oxidation or N-dearylation of the benzisothiazole moiety. Prakash, C. *et al.*, Drug Metab. Dispos., 25(7):863-872 (1997). At least 12 human metabolites have been identified: ziprasidone sulfoxide (ZIP-SO); ziprasidone sulfone (ZIP-SO₂); 3-(piperazine-1-yl)-1,2-benzisothiazole

(BITP); BITP sulfoxide; BITP sulfone; 6-chloro-5-(2-piperazin-1-yl-ethyl)-1,3-dihydro-indol-2-one; 6-chloro-5-(2-{4-[imino-(2-mercapto-phenyl)methyl]-piperazin-1-yl}ethyl)-1,3-dihydro-indol-2-one; 6-chloro-5-(2-{4-[imino-(2-methylsulfanyl-phenyl)methyl]-piperazin-1-yl}ethyl)-1,3-dihydro-indol-2-one; S-methyl-dihydro-ziprasidone; S-methyl-dihydro-ziprasidone sulfoxide; dihydro-ziprasidone sulfoxide; and (6-chloro-2-oxo-2,3-dihydro-1H-indol-5-yl)acetic acid. Two metabolites, ZIP-SO and ZIP-SO₂, both of which are formed by oxidation of the ziprasidone sulfur atom are discussed herein. These metabolites have the following structures:



Ziprasidone Sulfoxide (ZIP-SO)



Ziprasidone Sulfone (ZIP-SO₂)

Both ZIP-SO and ZIP-SO₂ are minor metabolites, and account for less than about 10% and less than about 3% of ziprasidone metabolites found in human urine, respectively. Prakash, C. *et al.*, Drug Metab. Dispos., 25(7):863-872 (1997). It has been reported that neither metabolite likely contributes to the antipsychotic activity of ziprasidone. Prakash, C. *et al.*, Drug Metab. Dispos., 25(7):863-872 (1997). Indeed, it has been reported that ziprasidone metabolites in general are not active at the D₂ and 5-HT_{2A} receptor sites. Ereshefsky, L., J. Clin. Psych., 57(suppl. 11):12-25 (1996).

Ziprasidone offers a number of benefits, but unfortunately many adverse effects are associated with its administration. Examples of adverse affects of ziprasidone include, but are not limited to, nausea, somnolence, asthenia, dizziness, extra-pyramidal symptoms, akathisia, cardiovascular disturbances, male sexual dysfunction, and elevated serum liver enzyme levels. Davis, R. and Markham, A., CNS Drugs, 8(2):154-159 (1997).

These adverse effects can significantly limit the dose level, frequency, and duration of drug therapy. It is thus desirable to find a compound which possesses advantages of ziprasidone but fewer of its disadvantages.

3. SUMMARY OF THE INVENTION

This invention relates to novel methods using, and compositions comprising, ziprasidone metabolites, preferably, ziprasidone sulfoxide and ziprasidone sulfone. These metabolites, prior to the present invention, have been reported to have little or no in vivo activity. The present invention encompasses the in vivo use of these metabolites, and their incorporation into pharmaceutical compositions and single unit dosage forms useful in the treatment and prevention of disorders that are ameliorated by the inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine reuptake at dopamine D₂ receptors. Such disorders include psychotic and neuroleptic disorders. In a preferred embodiment, ziprasidone metabolites are used in the treatment or prevention of neuroleptic and related disorders in mammals, including humans.

The compounds and compositions of the invention further allow the treatment and prevention of the diseases and disorders while reducing or avoiding adverse effects associated with the administration of ziprasidone.

3.1 DEFINITIONS

As used herein, the term "patient" refers to a mammal, particularly a human.

As used herein, the term "ziprasidone metabolite" means a compound that is a product of the metabolism of ziprasidone in a human. Ziprasidone metabolites include, but are not limited to: ziprasidone sulfoxide (ZIP-SO); ziprasidone sulfone (ZIP-SO₂); 3-(piperazine-1-yl)-1,2-benzisothiazole (BITP); BITP sulfoxide; BITP sulfone; 6-chloro-5-(2-piperazin-1-yl-ethyl)-1,3-dihydro-indol-2-one; 6-chloro-5-(2-{4-[imino-(2-mercapto-phenyl)methyl]-piperazin-1-yl}ethyl)-1,3-dihydro-indol-2-one; 6-chloro-5-(2-{4-[imino-(2-methylsulfanyl-phenyl)methyl]-piperazin-1-yl}ethyl)-1,3-dihydro-indol-2-one; S-methyl-dihydro-ziprasidone; S-methyl-dihydro-ziprasidone sulfoxide; dihydro-ziprasidone sulfoxide; and (6-chloro-2-oxo-2,3-dihydro-1H-indol-5-yl)acetic acid. Preferred ziprasidone metabolites include ZIP-SO and ZIP-SO₂.

As used herein, the term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic acids, including inorganic acids and organic acids. Suitable non-toxic acids include, but are not limited to, inorganic and organic acids such as acetic, alginic, anthranilic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethenesulfonic, formic, fumaric, furoic, galacturonic, gluconic,

glucuronic, glutamic, glycolic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phenylacetic, phosphoric, propionic, salicylic, stearic, succinic, sulfanilic, sulfuric, tartaric acid, and p-toluenesulfonic acid. Preferred non-toxic acids include hydrochloric, hydrobromic, phosphoric, sulfuric, and methanesulfonic acids. Examples of preferred salts thus include hydrochloride and mesylate salts.

As used herein, the term "a method of treating disorders ameliorated by the inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine reuptake at dopamine D₂ receptors in a patient" means relief from symptoms of disease states associated with abnormal serotonin and/or dopamine levels; such symptoms are reduced or relieved by way of inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine reuptake at dopamine D₂ receptors in a patient. Disorders treated by inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine reuptake at dopamine D₂ receptors in a patient include, but are not limited to, neuroleptic disorders, migraines, acute intermittent porphyria, intractable hiccups, Parkinson's disease and epilepsy.

As used herein, the term "psychosis" means a mental or behavioral disorder, with or without organic damage, causing gross distortion or disorganization of a person's mental capacity, affective response, capacity to recognize reality, communicate, or relate to others such that his or her capacity to cope with the ordinary demands of everyday life is diminished. Psychosis includes, but is not limited to, hallucinations, paranoia, affective psychosis (manic psychosis), alcoholic psychoses, arteriosclerotic psychosis, amnestic psychosis, bipolar psychosis (manic-depressive psychosis), Cheyne-Stokes psychosis, climacteric psychosis, depressive psychosis, drug psychosis, dysmnestic psychosis, hysterical psychosis, infection-exhaustion psychosis, Korsakoff's psychosis, postinfectious psychosis, postpartum psychosis, posttraumatic psychosis, senile psychosis, situational psychosis, toxic psychosis, traumatic psychosis, Windigo psychosis, schizo-affective psychosis, schizophrenia and related disorders. Diagnostic and Statistical Manual of Mental Disorders, 4th Ed., American Psychiatric Association (1997) (DSM-IV™)

As used herein, the term "affective disorder" means a disorder selected from the group including, but not limited to, depression, attention deficit disorder, attention deficit disorder with hyperactivity, and bipolar and manic conditions. The terms "attention deficit disorder" (ADD) and "attention deficit disorder with hyperactivity" (ADHD), or attention deficit/hyperactivity disorder (AD/HD), are used herein in accordance with the accepted meanings as found in the Diagnostic and Statistical Manual of Mental Disorders, 4th Ed., American Psychiatric Association (1997) (DSM-IV™), and Diagnostic and

Statistical Manual of Mental Disorders, 3rd Ed., American Psychiatric Association (1981) (DSM-IIITM).

As used herein, the term "a method of treating or preventing depression" means relief from the symptoms of depression which include, but are not limited to, changes in mood, feelings of intense sadness, despair, mental slowing, loss of concentration, pessimistic worry, agitation, and self-deprecation. Physical changes may also be relieved, including insomnia, anorexia, weight loss, decreased energy and libido, and abnormal hormonal circadian rhythms.

As used herein, the term "anxiety" is consistent with accepted meaning in the art. See, e.g., DSM-IVTM. Anxiety includes, but is not limited to, anxiety attacks, free-floating anxiety, noetic anxiety, separation anxiety, and situation anxiety. The terms "methods of treating or preventing" when used in connection with these disorders means amelioration, prevention or relief from the symptoms and/or effects associated with these disorders.

As used herein, the term "adverse effects of ziprasidone" means an effect selected from the group including, but not limited to, nausea, somnolence, asthenia, dizziness, motor disturbances (extrapyramidal symptoms), akathisia, cardiovascular disturbances (postural hypotension and tachycardia), respiratory disorder (described as coryzal symptoms, not nasal stuffiness), headache, dyspepsia, male sexual dysfunction, and elevated serum liver enzyme levels.

4. DETAILED DESCRIPTION OF THE INVENTION

The invention relates to methods of treating neuroleptic and related disorders using ziprasidone metabolites, and using ZIP-SO and ZIP-SO₂ in particular. Until now, ZIP-SO and ZIP-SO₂ were believed to possess little or no pharmacological activity. This invention further relates to solid and liquid pharmaceutical compositions and single unit dosage forms comprising a ziprasidone metabolite, such as ZIP-SO and ZIP-SO₂, as well as to methods of making ZIP-SO and ZIP-SO₂.

The methods and compositions of the invention can be used in the treatment and prevention of disorders described herein while avoiding or reducing drug-drug interactions and other adverse effects associated with agents known for the treatment of such disorders, including ziprasidone. The ziprasidone metabolites of the invention may further provide an overall improved therapeutic index over ziprasidone.

A first embodiment of the invention encompasses a method of treating or preventing disorders ameliorated by the inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine reuptake at dopamine D₂ receptors in a patient. The

5-HT₂ and D₂ receptors may be centrally (*i.e.*, in the central nervous system) or peripherally located. This method comprises administering to a patient in need of such treatment or prevention a therapeutically effective amount of a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof. Preferred ziprasidone metabolites include ZIP-SO and ZIP-SO₂. Disorders ameliorated by the inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine reuptake at dopamine D₂ receptors include, but are not limited to, neuroleptic disorders, pain, migraines, acute intermittent porphyria, intractable hiccups, Parkinson's disease and epilepsy. Neuroleptic disorders include, but are not limited to, psychosis, affective disorders, and anxiety.

A preferred embodiment of the invention thus encompasses a method of treating or preventing psychosis in a patient which comprises administering to a patient in need of such treatment or prevention a therapeutically effective amount of a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof. This embodiment encompasses methods of treating and preventing schizophrenia, schizoaffective psychosis, hallucinations, paranoia, affective psychosis (manic psychosis), alcoholic psychoses, arteriosclerotic psychosis, amnesic psychosis, bipolar psychosis (manic-depressive psychosis), Cheyne-Stokes psychosis, climacteric psychosis, depressive psychosis, drug psychosis, dysmnestic psychosis, hysterical psychosis, infection-exhaustion psychosis, Korsakoff's psychosis, postinfectious psychosis, postpartum psychosis, posttraumatic psychosis, senile psychosis, situational psychosis, toxic psychosis, traumatic psychosis, and Windigo psychosis.

Another preferred embodiment of the invention encompasses a method of treating or preventing an affective disorder in a patient which comprises administering to a patient in need of such treatment or prevention a therapeutically effective amount of a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof. This embodiment encompasses methods of treating and preventing depression, attention deficit disorder, attention deficit disorder with hyperactivity, combativeness, explosive hyperexcitable behavior, and bipolar and manic conditions.

A further preferred embodiment of the invention encompasses a method of treating and preventing anxiety in a patient which comprises administering to a patient in need of such treatment or prevention a therapeutically effective amount of a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof. This embodiment encompasses methods of treating and preventing anxiety attacks, free-floating anxiety, noetic anxiety, separation anxiety, and situation anxiety.

Another embodiment of the invention encompasses a method for treating and preventing pain in a patient which comprises administering to a patient in need of such treatment or prevention a therapeutically effective amount of a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof.

5 In a particular method encompassed by this embodiment, a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof, is adjunctively administered with at least one additional therapeutic agent. Examples of additional therapeutic agents include, but are not limited to: tricyclic antidepressants such as desipramine, imipramine, amitriptyline, and nortriptyline; anticonvulsants such as
10 carbamazepine and valproate; serotonin reuptake inhibitors such as fluoxetine, paroxetine, sertraline, and methysergide; mixed serotonin-norepinephrine reuptake inhibitors such as venlafaxine and duloxetine; serotonin receptor agonists; cholinergic (muscarinic and nicotinic) analgesics such as ketoprofen, aspirin, acetaminophen, indomethacin, ketorolac, and methotrimeprazine; adrenergic agents; neurokinin antagonists; xanthine oxidase
15 inhibitors such as allopurinol; and pharmaceutically acceptable salts and solvates thereof.

A second embodiment of the invention encompasses pharmaceutical compositions comprising a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof. Preferred ziprasidone metabolites include ZIP-SO and ZIP-SO₂. This embodiment further encompasses individual dosage forms of ziprasidone
20 metabolites, or pharmaceutically acceptable salts, solvates, hydrates, or clathrates thereof. Individual dosage forms of the invention may be suitable for oral, mucosal (including rectal, nasal, or vaginal), parenteral (including subcutaneous, intramuscular, bolus injection, intraarterial, or intravenous), sublingual, transdermal, buccal, or topical administration.

A particular pharmaceutical composition encompassed by this embodiment
25 comprises a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof, and at least one additional therapeutic agent. Examples of additional therapeutic agents include, but are not limited to: tricyclic antidepressants such as desipramine, imipramine, amitriptyline, and nortriptyline; anticonvulsants such as carbamazepine and valproate; serotonin reuptake inhibitors such as fluoxetine, paroxetine,
30 sertraline, and methysergide; mixed serotonin-norepinephrine reuptake inhibitors such as venlafaxine and duloxetine; serotonin receptor agonists; cholinergic (muscarinic and nicotinic) analgesics such as ketoprofen, aspirin, acetaminophen, indomethacin, ketorolac, and methotrimeprazine; adrenergic agents; neurokinin antagonists; xanthine oxidase inhibitors such as allopurinol; and pharmaceutically acceptable salts and solvates thereof.

35 A third embodiment of the invention encompasses methods of preparing ZIP-SO and ZIP-SO₂. These methods comprise treating ziprasidone with at least one

oxidizing agent. Preferably, the oxidizing agent is selected from the group consisting of hydrogen peroxide; sodium periodate; alkylperoxides; alkylhydroperoxides; hypochlorites, such as sodium hypochlorite and calcium hypochlorite; dioxiranes; nitric acid and a group VIII, IB and IIB transition metal catalyst; molecular oxygen or air and a lanthanide or transition metal catalyst; acyl nitrites; sodium perborate; and peracids.

4.1. SYNTHESIS AND PREPARATION

Ziprasidone sulfoxide (ZIP-SO) and ziprasidone sulfone (ZIP-SO₂) are readily prepared from ziprasidone using oxidation methods known to those skilled in the art. A syntheses of ziprasidone are described in U.S. Patent Nos. 4,831,031; 5,206,366; 5,338,846; and 5,359,068, the disclosure of which is hereby incorporated by reference.

In general, sulfoxides are formed by oxidation of thioalkyl groups using one mole equivalent of an oxidizing agent. Sulfoxides can be further oxidized to sulfones by using a second mole of an oxidizing agent. Preferably, the oxidizing agent is hydrogen peroxide; sodium periodate; alkylperoxides; alkylhydroperoxides; hypochlorites, such as sodium hypochlorite and calcium hypochlorite; dioxiranes; nitric acid and a gold tetrachloride catalyst; potassium permanganate; sodium perborate; potassium hydrogen persulfate; molecular oxygen and a ceric ammonium nitrate catalyst; acyl nitrites; sodium perborate; and peracids. March, J., Advanced Organic Chemistry, 4th Edition, John Wiley & Sons, pp.1201-1203 (1992). When sufficient amounts of oxidizing agent are present, thioalkyl groups can be converted directly to sulfones without isolation of sulfoxides. If necessary, the nitrogen of the benzisothialolyl ring can be protected using suitable methods known to those skilled in the art; an example is the reaction with anhydride to yield the corresponding amide, which can be removed after oxidation of sulfur. See, e.g., March, J. Advanced Organic Chemistry, 4th Edition p. 401 and 418-419 (1985). Suitable solvents include acetonitrile, methylene chloride, benzene, toluene, N-methylpyrrolidinone, dimethylformamide, ethanol, methanol, isopropanol, propanol, butanol, isobutanol, *tert*-butyl alcohol, dimethylsulfoxide, diethyl ether, tetrahydrofuran, acetone, and mixtures thereof, including aqueous mixtures where appropriate.

4.2. PHARMACEUTICAL COMPOSITIONS AND METHOD OF USE

The active compounds of the invention (*i.e.*, ziprasidone metabolites) are antipsychotic and antineuroleptic agents, and may thus be used in the treatment or prevention of a wide range of diseases and conditions. The magnitude of a prophylactic or therapeutic dose of a particular active ingredient of the invention in the acute or chronic management of a disease or condition will vary, however, with the nature and severity of

the disease or condition, and the route by which the active ingredient is administered. The dose, and perhaps the dose frequency, will also vary according to the age, body weight, and response of the individual patient. Suitable dosing regimens can be readily selected by those skilled in the art with due consideration of such factors. In general, the recommended daily dose range for the conditions described herein lie within the range of from about 1 mg to about 1000 mg per day, given as a single once-a-day dose in the morning but preferably as divided doses throughout the day taken with food. More preferably, the daily dose is administered twice daily in equally divided doses. Preferably, a daily dose range should be from about 5 mg to about 500 mg per day, more preferably, between about 10 mg and about 200 mg per day. In managing the patient, the therapy should be initiated at a lower dose, perhaps about 1 mg to about 25 mg, and increased if necessary up to about 200 mg to about 1000 mg per day as either a single dose or divided doses, depending on the patient's global response.

It may be necessary to use dosages of the active ingredient outside the ranges disclosed herein in some cases, as will be apparent to those of ordinary skill in the art. Because elimination of ziprasidone metabolites from the bloodstream is dependant on renal and liver function, it is recommended that the total daily dose be reduced by at least 50% in patients with moderate hepatic impairment, and that it be reduced by 25% in patients with mild to moderate renal impairment. For patients undergoing hemodialysis, it is recommended that the total daily dose be reduced by 5% and that the dose be withheld until the dialysis treatment is completed. Furthermore, it is noted that the clinician or treating physician will know how and when to interrupt, adjust, or terminate therapy in conjunction with individual patient response.

The phrase "therapeutically effective amount," as used herein with respect to the treatment or prevention of disorders ameliorated by the inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine reuptake at dopamine D₂ receptors, such as neuroleptic disorders, encompasses the above described dosage amounts and dose frequency schedules. Different therapeutically effective amounts may be applicable for different diseases and conditions, as will be readily known by those of ordinary skill in the art. Similarly, amounts sufficient to treat or prevent such disorders, but insufficient to cause, or sufficient to reduce, adverse effects associated with ziprasidone, are also encompassed by the above described dosage amounts and dose frequency schedules.

Any suitable route of administration may be employed for providing the patient with an effective dosage of a ziprasidone metabolite. For example, oral, mucosal (including rectal), parenteral (including subcutaneous, intramuscular, bolus injection, and intravenous), sublingual, transdermal, nasal, buccal, and like may be employed. In the

acute treatment or management of a disease or condition, it is preferred that the active ingredient be administered orally. In the acute treatment or management of a disease or condition, it is preferred that the active ingredient be administered parenterally.

The pharmaceutical compositions of the invention comprise at least one
5 ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof as an active ingredient, and may also contain a pharmaceutically acceptable carrier and optionally other therapeutic ingredients known to those skilled in the art, including the additional therapeutic ingredients listed above. The pharmaceutical compositions may be solid or liquid. Examples of solid compositions include crystalline, non-crystalline (i.e.,
10 amorphous), hydrated, and anhydrous compositions. Preferred pharmaceutical compositions are hydrates, including, but not limited to, mesylate dihydrates, mesylate trihydrates, and hydrochloride monohydrates. Such hydrates are described in U.S. Patent No. 5,312,925, PCT Publication No. WO/97/42190, and PCT Publication No. WO/97/42191, the disclosures of which are each incorporated herein. The pharmaceutical
15 compositions may also be inclusion complexes, such as those described in PCT Publication No. WO 97/41896, the disclosure of which is incorporated herein.

Compositions of the invention are suitable for oral, mucosal (including rectal), parenteral (including subcutaneous, intramuscular, bolus injection, and intravenous), sublingual, transdermal, nasal, or buccal administration, although the most suitable route in
20 any given case will depend on the nature and severity of the condition being treated. The compositions may be conveniently presented in unit dosage form and prepared by any of the methods well known in the part of pharmacy. Dosage forms include tablets, caplets, troches, lozenges, dispersions, suspensions, suppositories, solutions, capsules, soft elastic gelatin capsules, patches, and the like. Preferred dosage forms are suitable for oral
25 administration. Lyophilized dosage forms may be orally administered, or may be reconstituted to provide sterile, liquid dosage forms suitable for parenteral administration to a patient.

In practical use, a ziprasidone metabolite can be combined as the active ingredient in intimate admixture with a pharmaceutically acceptable carrier according to
30 conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms and comprises a number of components depending on the form of preparation desired for administration. The compositions of the invention include, but are not limited to, suspensions, solutions and elixirs; aerosols; or excipients, including, but not limited to, starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders,
35 disintegrating agents, and the like. Preferably, the pharmaceutical composition is in the form of an oral preparation.

Pharmaceutical compositions of the invention suitable for oral administration may be presented as discrete pharmaceutical unit dosage forms, such as capsules, cachets, soft elastic gelatin capsules, tablets, caplets, or aerosols sprays, each containing a predetermined amount of the active ingredients, as a powder or granules, or as a solution or
5 a suspension in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil liquid emulsion. Such compositions may be prepared by any method known in the art of pharmacy which comprises the step of bringing an active ingredient into association with a carrier. In general, the compositions are prepared by uniformly and intimately admixing the active ingredients with liquid carriers or finely divided solid
10 carriers or both, and then, if necessary, shaping the product into the desired presentation. Oral solid preparations are preferred over oral liquid preparations. Preferred oral solid preparations are capsules and tablets.

A tablet may be prepared by compression or molding techniques. Compressed tablets may be prepared by compressing in a suitable machine the active
15 ingredient in a free-flowing form, such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, granulating agent, surface active or dispersing agent, or the like. Molded tablets may be made by molding, in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Preferably, each tablet, cachet, caplet, or capsule contains from about 1 mg to about 1000 mg of ziprasidone metabolite,
20 more preferably from about 5 mg to about 500 mg, and most preferably from about 10 mg to about 200 mg.

Pharmaceutical compositions of the invention may also be formulated as a pharmaceutical composition in a soft elastic gelatin capsule unit dosage form by using conventional methods well known in the art. See, e.g., Ebert, Pharm. Tech., 1(5):44-50
25 (1977). Soft elastic gelatin capsules have a soft, globular gelatin shell somewhat thicker than that of hard gelatin capsules, wherein a gelatin is plasticized by the addition of plasticizing agent, e.g., glycerin, sorbitol, or a similar polyol. The hardness of the capsule shell may be changed by varying the type of gelatin used and the amounts of plasticizer and water. The soft gelatin shells may contain a preservative, such as methyl- and
30 propylparabens and sorbic acid, to prevent the growth of fungi. The active ingredient may be dissolved or suspended in a liquid vehicle or carrier, such as vegetable or mineral oils, glycols, such as polyethylene glycol and propylene glycol, triglycerides, surfactants, such as polysorbates, or a combination thereof.

A pharmaceutically acceptable excipient used in the compositions and
35 dosage form of the invention may be a binder, a filler, a mixture thereof. A pharmaceutically acceptable excipient may also include a lubricant, a disintegrant, or

mixtures thereof. Preferred excipients are lactose, croscarmellose, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate. One embodiment of the invention encompasses a pharmaceutical composition which is substantially free of all mono- or di-saccharide excipients.

5 Binders suitable for use in the compositions and dosage forms of the invention include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose),
10 polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208, 2906, 2910), microcrystalline cellulose or mixtures thereof.

Suitable forms of microcrystalline cellulose include, for example, the materials sold as AVICEL-PH-101, AVICEL-PH-103 and AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA.,
15 U.S.A.). An exemplary suitable binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581 by FMC Corporation.

Fillers suitable for use in the compositions and dosage forms of the invention include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid,
20 sorbitol, starch, pre-gelatinized starch, or mixtures thereof.

The binder/filler in pharmaceutical compositions of the invention is typically present in about 50 to about 99 weight percent of the pharmaceutical composition.

Disintegrants are used to cause the tablet to disintegrate when exposed to an aqueous environment. Too much of a disintegrant will produce tablets which may
25 disintegrate in the bottle due to atmospheric moisture; too little may be insufficient for disintegration to occur and may thus alter the rate and extent of release of the drug ingredient(s) from the dosage form. Thus, a sufficient amount of disintegrant that is neither too little nor too much to detrimentally alter the release of the drug ingredient(s) should be used to form dosage forms of ziprasidone metabolite made according to the invention. The
30 amount of disintegrant used varies based upon the type of formulation and mode of administration, and is readily discernible to those of ordinary skill in the art. Typically, about 0.5 to about 15 weight percent of disintegrant, preferably about 1 to about 5 weight percent of disintegrant, may be used in the pharmaceutical composition.

Disintegrants suitable for use in the compositions and dosage forms of the
35 invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium,

sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other algin, other celluloses, gums or mixtures thereof.

Lubricants suitable for use in the compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, 5 light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laurate, agar, or mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL 200, manufactured by W.R. Grace Co. of Baltimore MD), a 10 coagulated aerosol of synthetic silica (marketed by Deaussa Co. of Plano, Texas), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, Mass), or mixtures thereof. A lubricant may optionally be added, typically in an amount of less than about 1 weight percent of the pharmaceutical composition.

In addition to the common dosage forms set out above, the compounds of the 15 invention may also be administered by controlled release means or delivery devices that are well known to those of ordinary skill in the art, such as those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, the disclosures of which are each incorporated herein by express reference thereto. These pharmaceutical 20 compositions can be used to provide slow or controlled-release of one or more of the active ingredients therein using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or the like, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of 25 ordinary skill in the art, including those described herein, may be readily selected for use with the pharmaceutical compositions of the invention. Thus, single unit dosage forms suitable for oral administration, such as tablets, capsules, gelcaps, caplets, and the like, that are adapted for controlled-release are encompassed by the invention.

All controlled-release pharmaceutical products have a common goal of 30 improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations may include: 1) extended activity of the drug; 2) reduced dosage frequency; 3) increased 35 patient compliance; and 4) a lower peak plasma concentration of the drug. The latter advantage is significant because high peak plasma concentrations of some drugs can cause

adverse effects not associated with lower, but still therapeutically effective, plasma concentrations.

Most controlled-release formulations are designed to initially release an amount of drug that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body.

The controlled-release of an active ingredient may be stimulated by various inducers, for example pH, temperature, enzymes, water, or other physiological conditions or compounds. The term "controlled-release component" in the context of the invention is defined herein as a compound or compounds, including, but not limited to, polymers, polymer matrices, gels, permeable membranes, liposomes, microspheres, or the like, or a combination thereof, that facilitates the controlled-release of the active ingredient.

Pharmaceutical compositions of the invention may also be formulated for parenteral administration by injection (subcutaneous, bolus injection, intramuscular, or intravenous), and may be dispensed in a unit dosage form, such as a multidose container or an ampule. Such compositions for parenteral administration may be in the form of suspensions, solutions, emulsions, or the like in aqueous or oily vehicles, and in addition to the active ingredients may contain one or more formulary agents, such as dispersing agents, suspending agents, stabilizing agents, preservatives, and the like.

The invention is further defined by reference to the following examples describing in detail the preparation of the compositions of the invention. It will be apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the purpose and interest of this invention.

5. EXAMPLES

5.1. EXAMPLE 1: SYNTHESIS OF ZIPRASIDONE

To a 125 mL round-bottom flask equipped with an N₂ inlet and condenser are added 0.73 g (3.2 mmol) 5-(2-chloroethyl)-6-chloro-oxindole, 0.70 g (3.2 mmol) N-(1,2-benzisothiazol-3-yl)piperazine, 0.68 g (6.4 mmol) sodium carbonate, 2 mg sodium iodide, and 30 mL methylisobutyl ketone. The reaction is refluxed for 40 hours, cooled, filtered, and evaporated. The residue is chromatographed on silica gel, eluting the by-products with ethyl acetate (1 L) and the product with 4 % methanol in ethyl acetate (1.5 L). The product fractions (R_f = 0.2 in 5 % methanol in ethyl acetate) are evaporated, taken up in methylene chloride, and precipitated by addition of ether saturated with HCl; the solid is filtered and

washed with ether, dried, and washed with acetone. The latter is done by slurrying the solid with acetone and filtering. Ziprasidone is obtained as a high melting, non-hygroscopic solid product having an expected melting point of 288°C to 288.5°C.

5 **5.2. EXAMPLE 2: SYNTHESIS OF ZIPRASIDONE SULFOXIDE**

To a solution of ziprasidone made as described in Example 1 (0.70 g, 1.7 mmol) in acetonitrile is added 30 % H₂O₂ (1.7 mmol). After stirring for 24 hours at room temperature, the reaction mixture is cooled, filtered, and evaporated. The residue is chromatographed on silica gel, eluting the by-products with ethyl acetate (1 L) and the
10 product with 4 % methanol in ethyl acetate (1.5 L). The product fractions are evaporated, taken up in methylene chloride, and precipitated by addition of ether saturated with HCl; the solid is filtered and washed with ether, dried, and washed with acetone.

5.3. EXAMPLE 3: SYNTHESIS OF ZIPRASIDONE SULFONE

15 To a solution of ziprasidone sulfoxide made as described in Example 2 (0.76 g, 1.7 mmol) in acetonitrile is added 30 % H₂O₂ (1.7 mmol). After stirring for 24 hours at room temperature, the reaction mixture is cooled, filtered, and evaporated. The residue is chromatographed on silica gel, eluting the by-products with ethyl acetate (1 L) and the product with 4 % methanol in ethyl acetate (1.5 L). The product fractions are evaporated,
20 taken up in methylene chloride, and precipitated by addition of ether saturated with HCl; the solid is filtered and washed with ether, dried, and washed with acetone.

Alternatively, ziprasidone sulfone may be obtained by one step oxidation of ziprasidone. To a solution of ziprasidone made as described in Example 1 (0.70 g, 1.7 mmol) in acetonitrile is added 30 % H₂O₂ (3.4 mmol). After stirring for 24 hours at room
25 temperature, the reaction mixture is cooled, filtered, and evaporated. The residue is chromatographed on silica gel, eluting the by-products with ethyl acetate (1 L) and the product with 4 % methanol in ethyl acetate (1.5 L). The product fractions are evaporated, taken up in methylene chloride, and precipitated by addition of ether saturated with HCl; the solid is filtered and washed with ether, dried, and washed with acetone.

30

5.4. EXAMPLE 4: 5-HT₂ RECEPTOR ACTIVITY

Receptor selection and amplification technology (R-SAT) is used (Receptor Technologies Inc., Winooski, VT) to determine potential agonist and/or antagonist activity of ziprasidone and ziprasidone metabolites on cloned human serotonin 5-HT₂ receptor
35 subtypes expressed in NIH 3T3 cells. This assay is a modification of a known assay to determine potential agonist and/or antagonist activity of racemic norcisapride, cisapride and

their enantiomers. (Burstein *et al.*, J. Biol Chem., 270:3141-3146 (1995); and Messier *et al.*, Pharmacol. Toxicol., 76(5):308-311 (1995)).

The assay involves co-expression of a marker enzyme, β -galactosidase, with the serotonin receptor of interest. Ligands stimulate proliferation of cells that express the
5 receptor and, therefore, the marker. Ligand-induced effects can be determined by assay of the marker.

NIH 3T3 cells are incubated, plated, and then transfected using human 5-HT₂ serotonin receptors, pSV- β -galactosidase, and salmon sperm DNA. The medium is changed one day later, and after 2 days, aliquots of the trypsinized cells are placed in wells of a 96
10 well plate. After five days in culture in the presence of the ligands, the levels of β -galactosidase are measured. The cells are then rinsed and incubated with the substrate, *o*-nitrophenyl β -D-galactopyranoside. After 16 hours, the plates are read at 405 nm on a plate-reader. Each compound is tested for activity in triplicate at seven different concentrations (10, 2.5, 0.625, 0.156, 0.039, 0.0098, and 0.0024 nM).

15.

5.5. EXAMPLE 5: DOPAMINE D₂ RECEPTOR ACTIVITY

Competition radioreceptor assays are used to determine the affinity (IC₅₀'s) of the phenylaminotetralins and other reference ligands for D₂ dopamine receptors. D₂ assays uses a 90 minute incubation with [³H]YM-09151-2 (0.065 nM) with (+)-butaclamol
20 (0.25 μ M) defining nonspecific binding. Jarvie, J.R. *et al.*, Eur. J. Pharmacol., 144:163-171 (1987) and Kula, N.S. *et al.*, Dev. Brain Res., 66:286-287 (1992). Under these conditions, the K_D of [³H]SCH23390 is 0.34 nM and that of [³H]-YM-09151-2 is 0.045 nM. Test agents are evaluated by running, in duplicate, six or more concentrations that bracketed the IC₅₀. Three replications are performed, and the resulting data are analyzed using the
25 ALLFIT program.

The binding of the novel radioligand [³H](\pm)-4 to brain membranes is characterized using assay conditions similar to those developed for the σ ligand [³H]DTG. Weber, E. *et al.*, Proc. Nat. Acad. Sci. U.S.A., 83:8784-8788 (1986). Briefly, frozen guinea pig brain (minus cerebellum; obtained from Keystone Biologicals, Cleveland, OH) is
30 thawed and homogenized (10 mL/g tissue) in ice-cold 10 mM Tris-HCl buffer containing 0.32 M sucrose, pH 7.0; the homogenate is centrifuged at 1000g for 15 minutes at 4°C and the supernatant recentrifuged at 31,000g for 15 minutes at 4°C. The P₂ pellet is suspended in 10 mM Tris buffer (pH 7.4, 25°C) at 3 mL/g tissue and incubated at room temperature for 15 min at 4°C. The resulting pellet is stored at -70°C in 10mM Tris (pH 7.4) at
35 20 mg protein/mL. To determine binding parameters, a ligand saturation curve is constructed with 1.0 mg of brain protein (50 μ L) in glass tubes (triplicate) containing six

concentrations (0.02-2.0 nM) of free ligand (F) in 50 mM Tris-HCl buffer, pH 7.4 (2.0 mL total volume), with excess BMY-14802 (5.0 μ M) used to define specific binding. Tubes are incubated for 60 minutes at 30°C and then filtered in a Brandel cell harvester through glass fiber sheets, subsequently cut and counted for tritium by liquid scintillation spectrometry.

- 5 Results first are plotted in Scatchard-Rosenthal linearized form as ratio of bound/free ligand (B/F) vs. specific binding (B), to provide estimates of apparent affinity K_D (slope) and binding site density B_{max} (x intercept); these values are verified with the LIGAND curve-fitting program adapted to the MacIntosh microcomputer. Munson, P.J. *et al.*, Analyt. Biochem., 107:220-239 (1980). Under these conditions, the K_D of [3 H]4 is 0.031 nM. For
- 10 competitive binding assays, tubes are incubated (60 min, 30°C) with 50 pM (*ca.* K_D) [3 H]4, with 5 μ M BMY-14802 used to define nonspecific binding. From 4-8 concentrations (10 pM to 10 μ M) of test compounds are used, and the resulting competition data are computer curve-fitted to determine $IC_{50} \pm SEM$.

15 5.6. HARD GELATIN CAPSULE DOSAGE FORMS

Table I provides the ingredients of suitable capsule forms of the pharmaceutical compositions of this invention.

TABLE I

20	Component	25 mg capsule	50 mg capsule	100 mg capsule
	Ziprasidone Sulfoxide	25	50	100
	Microcrystalline Cellulose	90.0	90.0	90.0
25	Pre-gelatinized Starch	100.3	97.8	82.8
	Croscarmellose	7.0	7.0	7.0
	Magnesium Stearate	0.2	0.2	0.2

- 30 The active ingredient (*i.e.*, ziprasidone sulfoxide) is sieved and blended with the excipients listed. The mixture is filled into suitably sized two-piece hard gelatin capsules using suitable machinery and methods well known in the art. See Remington's Pharmaceutical Sciences, 16th or 18th Editions, each incorporated herein in its entirety by reference thereto. Other doses may be prepared by altering the fill weight and, if necessary,
- 35 by changing the capsule size to suit. Any of the stable hard gelatin capsule formulations above may be formed.

5.7. HARD GELATIN CAPSULE DOSAGE FORMS

Table II provides the ingredients of suitable capsule forms of the pharmaceutical compositions of this invention.

TABLE II

Component	25 mg capsule	50 mg capsule	100 mg capsule
Ziprasidone Sulfone	25	50	100
Microcrystalline Cellulose	90.0	90.0	90.0
Pre-gelatinized Starch	100.3	97.8	82.8
Croscarmellose	7.0	7.0	7.0
Magnesium Stearate	0.2	0.2	0.2

The active ingredient (i.e., ziprasidone sulfone) is sieved and blended with the excipients listed. The mixture is filled into suitably sized two-piece hard gelatin capsules using suitable machinery and methods well known in the art. Other doses may be prepared by altering the fill weight and, if necessary, by changing the capsule size to suit. Any of the stable hard gelatin capsule formulations above may be formed.

5.8. COMPRESSED TABLET DOSAGE FORMS

The ingredients of compressed tablet forms of the pharmaceutical compositions of the invention are provided in Table III.

TABLE III

Component	25 mg capsule	50 mg capsule	100 mg capsule
Ziprasidone Sulfoxide	25	50	100
Microcrystalline Cellulose	90.0	90.0	90.0
Pre-gelatinized Starch	100.3	97.8	82.8
Croscarmellose	7.0	7.0	7.0
Magnesium Stearate	0.2	0.2	0.2

The active ingredient (i.e., ziprasidone sulfoxide) is sieved through a suitable sieve and blended with the excipients until a uniform blend is formed. The dry blend is screened and blended with the magnesium stearate. The resulting powder blend is then compressed into tablets of desired shape and size. Tablets of other strengths may be prepared by altering the ratio of the active ingredient to the excipient(s) or modifying the table weight.

While the invention has been described with respect to the particular embodiments, it will be apparent to those skilled in the art that various changes and modifications may be made without departing from the spirit and scope of the invention as defined in the claims.

Such modifications are also intended to fall within the scope of the appended claims.

15

20

25

30

35

THE CLAIMS

What is claimed is:

- 5 1. A method of treating or preventing a disorder ameliorated by the inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine reuptake at dopamine D₂ receptors in a patient which comprises administering to a patient in need of such treatment or prevention a therapeutically effective amount of a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof.
- 10 2. The method of claim 1 wherein the disorder is selected from the group consisting of neuroleptic disorders, migraines, acute intermittent porphyria, intractable hiccups, Parkinson's disease and epilepsy.
- 15 3. A method of treating or preventing a neuroleptic disorder in a patient which comprises administering to a patient in need of such treatment or prevention a therapeutically effective amount of a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof.
- 20 4. The method of claim 1 or 3 wherein the patient is a human.
5. The method of claim 1 or 3 wherein the ziprasidone metabolite is ziprasidone sulfoxide or ziprasidone sulfone.
- 25 6. The method of claim 3 wherein the neuroleptic disorder is selected from the group consisting of psychosis, affective disorders, and anxiety.
7. The method of claim 6 wherein the psychosis is selected from the group consisting of schizophrenia, schizo-affective psychosis, hallucinations, paranoia, affective psychosis, alcoholic psychoses, arteriosclerotic psychosis, amnestic psychosis, bipolar psychosis, Cheyne-Stokes psychosis, climacteric psychosis, depressive psychosis, drug psychosis, dysmnesic psychosis, hysterical psychosis, infection-exhaustion psychosis, Korsakoff's psychosis, postinfectious psychosis, postpartum psychosis, posttraumatic psychosis, senile psychosis, situational psychosis, toxic psychosis, traumatic psychosis, and
30 Windigo psychosis.
35

8. The method of claim 6 wherein the affective disorder is selected from the group consisting of depression, attention deficit disorder, attention deficit disorder with hyperactivity, bipolar conditions and manic conditions.

5 9. The method of claim 6 wherein the anxiety is selected from the group consisting of anxiety attacks, free-floating anxiety, noetic anxiety, separation anxiety, and situation anxiety.

10 10. The method of claim 1 or 3 wherein the ziprasidone metabolite is administered parenterally, transdermally, mucosally, nasally, buccally, sublingually, or orally.

11. The method of claim 10 wherein the ziprasidone metabolite is administered orally.

15 12. The method of claim 11 wherein the ziprasidone metabolite administered orally in a tablet or capsule form.

13. The method of claim 1 or 3 wherein the therapeutically effective
20 amount of ziprasidone metabolite is between about 1 mg and about 1000 mg per day.

14. The method of claim 13 wherein the therapeutically effective amount of ziprasidone metabolite is between about 5 mg to about 500 mg per day.

25 15. The method of claim 14 wherein therapeutically effective amount of ziprasidone metabolite is between about 10 mg to about 200 mg per day.

16. A pharmaceutical composition comprising a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof.

30 17. The pharmaceutical composition of claim 16 wherein the ziprasidone metabolite is ziprasidone sulfoxide or ziprasidone sulfone.

18. The pharmaceutical composition of claim 16 wherein said
35 pharmaceutical composition further comprises an additional therapeutic agent selected from the group consisting of: tricyclic antidepressants; anticonvulsants; serotonin reuptake

inhibitors; mixed serotonin-norepinephrine reuptake inhibitors; serotonin receptor agonists; cholinergic analgesics; adrenergic agents; neurokinin antagonists; xanthine oxidase inhibitors; and pharmaceutically acceptable salts and solvates thereof.

5 19. The pharmaceutical composition of claim 18 wherein the tricyclic antidepressant is selected from the group consisting of desipramine, imipramine, amitriptyline, and nortriptyline.

 20. The pharmaceutical composition of claim 18 wherein the
10 anticonvulsant is selected from the group consisting of carbamazepine and valproate.

 21. The pharmaceutical composition of claim 18 wherein the serotonin reuptake inhibitor is selected from the group consisting of fluoxetine, paroxetine, sertraline, and methysergide.

15 22. The pharmaceutical composition of claim 18 wherein the mixed serotonin reuptake inhibitor is selected from the group consisting of venlafaxine and duloxetine.

20 23. The pharmaceutical composition of claim 18 wherein the cholinergic analgesic is selected from the group consisting of ketoprofen, aspirin, acetaminophen, indomethacin, ketorolac, and methotrimeprazine.

 24. The pharmaceutical composition of claim 18 wherein the xanthine
25 oxidase inhibitor is allopurinol.

 25. The pharmaceutical composition of claim 16 wherein said pharmaceutical composition further comprises a pharmaceutically acceptable carrier.

30 26. The pharmaceutical composition of claim 16 wherein said pharmaceutical composition is suitable for parenteral, transdermal, mucosal, nasal, buccal, sublingual, or oral administration to a patient.

 27. The pharmaceutical composition of claim 26 wherein said
35 pharmaceutical composition is suitable for oral administration to a patient.

28. The pharmaceutical composition of claim 16 wherein the amount of ziprasidone metabolite is between about 1 mg and about 1000 mg.

29. The pharmaceutical composition of claim 28 wherein the amount of ziprasidone metabolite is between about 5 mg and about 500 mg.

30. The pharmaceutical composition of claim 29 wherein the amount of ziprasidone metabolite is between about 10 mg and about 200 mg per day.

31. A dosage form suitable for the treatment and prevention of a neuroleptic disorder or pain which comprises a therapeutically effective amount of a ziprasidone metabolite, or a pharmaceutically acceptable salt, solvate, hydrate, or clathrate thereof.

32. The dosage form of claim 31 wherein the ziprasidone metabolite is ziprasidone sulfoxide or ziprasidone sulfone.

33. The dosage form of claim 31 wherein said pharmaceutical composition further comprises an additional therapeutic agent selected from the group consisting of: tricyclic antidepressants; anticonvulsants; serotonin reuptake inhibitors; mixed serotonin-norepinephrine reuptake inhibitors; serotonin receptor agonists; cholinergic analgesics; adrenergic agents; neurokinin antagonists; xanthine oxidase inhibitors; and pharmaceutically acceptable salts and solvates thereof.

34. The dosage form of claim 33 wherein the tricyclic antidepressant is selected from the group consisting of desipramine, imipramine, amitriptyline, and nortriptyline.

35. The dosage form of claim 33 wherein the anticonvulsant is selected from the group consisting of carbamazepine and valproate.

36. The dosage form of claim 33 wherein the serotonin reuptake inhibitor is selected from the group consisting of fluoxetine, paroxetine, sertraline, and methysergide.

37. The dosage form of claim 33 wherein the mixed serotonin reuptake inhibitor is selected from the group consisting of venlafaxine and duloxetine.
38. The dosage form of claim 33 wherein the cholinergic analgesic is
5 selected from the group consisting of ketoprofen, aspirin, acetaminophen, indomethacin, ketorolac, and methotrimeprazine.
39. The dosage form of claim 33 wherein the xanthine oxidase inhibitor
10 is allopurinol.
40. The dosage form of claim 31 wherein said dosage form further
comprises a pharmaceutically acceptable carrier.
41. The dosage form of claim 31 wherein said dosage form is suitable for
15 parenteral, transdermal, mucosal, nasal, buccal, sublingual, or oral administration to a patient.
42. The dosage form of claim 41 wherein said dosage form is a capsule
or a tablet.
20
43. The dosage form of claim 31 wherein the amount of ziprasidone
metabolite is between about 1 mg and about 1000 mg.
44. The dosage form of claim 43 wherein the amount of ziprasidone
25 metabolite is between about 5 mg and about 500 mg.
45. The dosage form of claim 44 wherein the amount of ziprasidone
metabolite is between about 10 mg and about 200 mg per day.
46. A method of preparing ziprasidone sulfoxide which comprises
30 treating ziprasidone with one molar equivalent of an oxidizing agent.
47. The method of claim 46 wherein the oxidizing agent is selected from
the group consisting of hydrogen peroxide; sodium periodate; alkylperoxides;
35 alkylhydroperoxides; hypochlorites, such as sodium hypochlorite and calcium hypochlorite; dioxiranes; nitric acid and a gold tetrachloride catalyst; potassium permanganate; sodium

perborate; potassium hydrogen persulfate; molecular oxygen and a ceric ammonium nitrate catalyst; acyl nitrites; sodium perborate; and peracids.

48. A method of preparing ziprasidone sulfone which comprises treating
5 ziprasidone with two molar equivalents of an oxidizing agent.

49. The method of claim 48 wherein the oxidizing agent is selected from
the group consisting of hydrogen peroxide; sodium periodate; alkylperoxides;
alkylhydroperoxides; hypochlorites, such as sodium hypochlorite and calcium hypochlorite;
10 dioxiranes; nitric acid and a gold tetrachloride catalyst; potassium permanganate; sodium
perborate; potassium hydrogen persulfate; molecular oxygen and a ceric ammonium nitrate
catalyst; acyl nitrites; sodium perborate; and peracids.

15

THIS PAGE BLANK (OPTIONAL)

20

25

30

35

THIS PAGE BLANK (USPTO)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 October 2000 (12.10.2000)

PCT

(10) International Publication Number
WO 00/59489 A3

- (51) International Patent Classification⁷: **A61K 31/495**
- (21) International Application Number: **PCT/US00/08707**
- (22) International Filing Date: **31 March 2000 (31.03.2000)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/127,939 **6 April 1999 (06.04.1999)** **US**
- (71) Applicant: **SEPRACOR INC. [US/US]; 111 Locke Drive, Marlborough, MA 01752 (US).**
- (72) Inventors: **BARBERICH, Timothy, J.; 40 Elm Street, Concord, MA 01742 (US). RUBIN, Paul, D.; 37 Greystone Lane, Sudbury, MA 01776 (US). YELLE, William, E.; 20 Ernie's Drive, Littleton, MA 01460 (US).**
- (74) Agents: **INSOGNA, Anthony, M. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).**
- (81) Designated States (*national*): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (*regional*): **ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**
- Published:**
— *With international search report.*
- (88) Date of publication of the international search report:
25 May 2001
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **METHODS AND COMPOSITIONS FOR THE TREATMENT OF PSYCHOTIC AND RELATED DISORDERS USING ZIPRASIDONE METABOLITES**

(57) Abstract: **The invention relates to novel methods using, and pharmaceutical compositions comprising, ziprasidone metabolites, preferably its sulfone and sulfoxide. The methods and compositions of the invention are suitable for the treatment of psychotic and related disorders and Parkinson's, discose, migraines, porpheria, epilepsy. The invention further encompasses methods of preparing ziprasidone sulfoxide and ziprasidone sulfone.**

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/08707

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/495

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, CHEM ABS Data, WPI Data, EMBASE, BIOSIS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PRAKASH, CHANDRA ET AL: "Metabolism and excretion of a new antipsychotic drug, ziprasidone, in humans" DRUG METAB. DISPOS. (1997), 25(7), 863-872	1-44
A	XP000987000 cited in the application the whole document ----- -/-	46-49

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

1 March 2001

Date of mailing of the international search report

09/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

A. Jakobs

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/08707

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PRAKASH, CHANDRA ET AL: "Metabolism and excretion of the novel antipsychotic drug ziprasidone in rats after oral administration of a mixture of 14C- and 3H-labeled ziprasidone" DRUG METAB. DISPOS. (1997), 25(2), 206-218 XP000987001 cited in the application	1-45
A	the whole document	46-49
X	EP 0 790 236 A (PFIZER) 20 August 1997 (1997-08-20) the whole document	1-4, 6-16, 18-31, 33-45
X	US 4 590 196 A (SMITH DAVID W ET AL) 20 May 1986 (1986-05-20) the whole document	16, 18-31, 33-45
X	YEVICH J P ET AL: "SYNTHESIS AND BIOLOGICAL EVALUATION OF 1-(1,2-BENZISOTHIAZOL-3-YL)-AND (1,2-BENZISOXAZOL-3-YL)PIPERAZINE DERIVATIVES AS POTENTIAL ANTIPSYCHOTIC AGENTS" JOURNAL OF MEDICINAL CHEMISTRY, US, AMERICAN CHEMICAL SOCIETY, WASHINGTON, vol. 29, no. 3, 1 March 1986 (1986-03-01), pages 359-369, XP000561328 ISSN: 0022-2623 the whole document	1-4, 6-16, 18-31, 33-45
X	US 4 831 031 A (LOWE III JOHN A ET AL) 16 May 1989 (1989-05-16) abstract column 1, line 24 -column 2, line 24	1-4, 6-16, 18-31, 33-45
X	WO 93 16073 A (WELLCOME FOUND) 19 August 1993 (1993-08-19) page 5, paragraph 7 page 50, paragraph 3 -page 51, paragraph 1; example 77	1-45
A	US 4 411 901 A (TEMPLE JR DAVIS L ET AL) 25 October 1983 (1983-10-25) the whole document	1

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Present claims 1,10-15 relate to a compound/therapeutic use defined (inter alia)

by reference to the following parameter(s):

P1: a disorder ameliorated by the inhibition of serotonin reuptake at 5-HT₂ receptors and/or the inhibition of dopamine D₂ receptors.

The use of these parameters in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT. It is impossible to compare the parameters the applicant has chosen to employ with what is set out in the prior art.

Re Claims 1-4,6-16,18-31,33-45: The term "ziprasidone metabolite" is not clear in the present context since it is not entirely clear into which compounds ziprasidone is metabolised by or in the human body.

Consequently, the search has been carried out for those parts of the application which do appear to be clear (and concise), namely the use of ziprasidone sulfoxide and ziprasidone sulfone in relation to the disorders specified in claims 2,3,6-9, and pharmaceutical compositions comprising ziprasidone sulfoxide and ziprasidone sulfone.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/08707

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0790236	A	20-08-1997	CA 2197272 A	14-08-1997
			JP 2911032 B	23-06-1999
			JP 9227516 A	02-09-1997
			US 5935960 A	10-08-1999
US 4590196	A	20-05-1986	AT 396685 B	25-11-1993
			AT 246985 A	15-03-1993
			AU 586182 B	06-07-1989
			AU 4653385 A	27-02-1986
			BE 903106 A	24-02-1986
			CA 1249587 A	31-01-1989
			CH 667268 A	30-09-1988
			DE 3530089 A	06-03-1986
			DK 381485 A	24-02-1986
			ES 546347 D	01-04-1987
			ES 8704485 A	16-06-1987
			FI 853193 A, B,	24-02-1986
			FR 2569404 A	28-02-1986
			GB 2163432 A, B	26-02-1986
			GR 852017 A	24-12-1985
			IE 58608 B	20-10-1993
			IT 1187730 B	23-12-1987
			JP 61112063 A	30-05-1986
			KR 9208820 B	09-10-1992
			LU 86051 A	28-02-1986
			NL 8502310 A	17-03-1986
			PT 81007 A, B	01-09-1985
			SE 464304 B	08-04-1991
			SE 8503922 A	24-02-1986
			ZA 8506387 A	30-04-1986
US 4831031	A	16-05-1989	US 4883795 A	28-11-1989
WO 9316073	A	19-08-1993	AU 3460393 A	03-09-1993
			CA 2117434 A	19-08-1993
			CZ 9401949 A	13-09-1995
			EP 0625978 A	30-11-1994
			FI 943718 A	11-08-1994
			HU 72309 A	29-04-1996
			JP 7503723 T	20-04-1995
			MX 9300786 A	31-03-1994
			NO 942977 A	10-10-1994
			SK 96494 A	11-07-1995
			ZA 9300958 A	11-08-1994
US 4411901	A	25-10-1983	AT 383124 B	25-05-1987
			AT 467782 A	15-10-1986
			AU 550780 B	10-04-1986
			AU 9087282 A	23-06-1983
			BE 895469 A	23-06-1983
			CA 1205806 A	10-06-1986
			CA 1216586 C	13-01-1987
			CH 659070 A	31-12-1986
			CH 656617 A	15-07-1986
			CY 1376 A	07-08-1987
			DE 3247530 A	30-06-1983
			DK 279183 A, B,	24-06-1983
			DK 307590 A, B,	28-12-1990

INTERNATIONAL SEARCH REPORT

Information on patent family members

In. tional Application No

PCT/US 00/08707

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4411901 A		DK 566282 A, B,	24-06-1983
		ES 518168 D	16-01-1984
		ES 8401970 A	01-04-1984
		ES 524946 D	01-04-1985
		ES 8503683 A	16-06-1985
		ES 525470 D	01-02-1985
		ES 8502991 A	01-05-1985
		ES 525471 D	01-02-1985
		ES 8503349 A	01-06-1985
		FI 824379 A, B,	24-06-1983
		FR 2521561 A	19-08-1983
		FR 2531431 A	10-02-1984
		GB 2114119 A, B	17-08-1983
		GR 77108 A	06-09-1984
		HK 5088 A	29-01-1988
		HU 195806 B	28-07-1988
		HU 190997 B	28-12-1986
		IE 54789 B	14-02-1990
		IT 1158037 B	18-02-1987
		JP 58110576 A	01-07-1983
		JP 1976527 C	17-10-1995
		JP 5017457 A	26-01-1993
		JP 5077669 B	27-10-1993
		KE 3718 A	16-04-1987
		KR 8701022 B	23-05-1987
		KR 8801378 B	29-07-1988
		LU 84550 A	08-09-1983
		NL 8204910 A	18-07-1983
		PT 76017 A, B	01-01-1983
		SE 453502 B	08-02-1988
		SE 8207350 A	10-08-1983
		SE 462162 B	14-05-1990
		SE 8702389 A	09-06-1987
		SG 30087 G	17-07-1987
		US 4452799 A	05-06-1984
		ZA 8208544 A	28-09-1983

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 October 2000 (12.10.2000)

PCT

(10) International Publication Number
WO 00/59489 A3

(51) International Patent Classification⁷: A61K 31/495

(21) International Application Number: PCT/US00/08707

(22) International Filing Date: 31 March 2000 (31.03.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/127,939 6 April 1999 (06.04.1999) US

(71) Applicant: SEPRACOR INC. [US/US]; 111 Locke Drive, Marlborough, MA 01752 (US).

(72) Inventors: BARBERICH, Timothy, J.; 40 Elm Street, Concord, MA 01742 (US); RUBIN, Paul, D.; 37 Greystone Lane, Sudbury, MA 01776 (US); YELLE, William, E.; 20 Ernie's Drive, Littleton, MA 01460 (US).

(74) Agents: INSOGNA, Anthony, M. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,

ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

(88) Date of publication of the international search report:
25 May 2001

(48) Date of publication of this corrected version:
27 September 2001

(15) Information about Correction:
see PCT Gazette No. 39/2001 of 27 September 2001, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND COMPOSITIONS FOR THE TREATMENT OF PSYCHOTIC AND RELATED DISORDERS USING ZIPRASIDONE METABOLITES

(57) Abstract: The invention relates to novel methods using, and pharmaceutical compositions comprising, ziprasidone metabolites, preferably its sulfone and sulfoxide. The methods and compositions of the invention are suitable for the treatment of psychotic and related disorders and Parkinson's disease, migraines, porphyria, epilepsy. The invention further encompasses methods of preparing ziprasidone sulfoxide and ziprasidone sulfone.

WO 00/59489 A3

THIS PAGE BLANK (USPTO)